contain target location information and the target would appear at either the left or right visual eld with equal possibility. Importantly, using the same approach as we used before (Huang et al., 2018; Jia et al., 2017; Liu et al., 2017), the luminance of the two discs, in both cueing and neutral conditions, was independently modulated for 5 s, according to two 5 s random temporal sequence respectively (Fig. 1C). We next employed the TRF approach (Crosse et al., 2016) to calculate and separate neural impulse response for the two discs (Att TRF and Unatt TRF) from the same EEG recordings. The TRF method, which uses linear regression to quantify the linear part of the stimulus-response relationships, models the neural impulse response function (i.e., the evoked brain response to a unit change in luminance in a stimulus sequence as a function of time lag).

As illustrated in Fig. 1C, we rst calculated the TRF responses for the attended, unattended, and neutral conditions in each channel and in each participant respectively (see representative participant data in Fig. 2A). We next performed a spectrotemporal analysis on the TRFs to examine their ne dynamic structures as a function of frequency (1–30 Hz) and time lag (0–0.8 s); this was done for each condition (Att, Unatt, Neutral), in each channel, and in each participant separately. Further analysis was then based on the spectrotemporal power pattern, in particular, the alpha-band power temporal pro le of the TRF responses for each condition, based on previous ndings supporting the critical role of alpha-band activity in visual attention (Haegens et al., 2012; Jia et al., 2017; Klimesch et al., 2007). Notably, the TRF spectrotemporal power difference between conditions might be not obvious in the temporal waveform since the grand average could cancel out the phase-inconsistent activities between subjects (Jia et al., 2017).

Fig. 2A illustrates the TRF responses of a representative participant on a representative sensor (Pz), and it is notable the TRF response became at and noisy when the relationship between the stimulus sequence and the corresponding trial response was shuf ed (achromatic lines), supporting that the calculated TRF waveform represented a genuine stimulus-speci c tracking response (Fig. 2A, left; see the reconstruction in terms of the TRFacti8dsponse andipt7

stimulus-speci

the Att Neutral comparison showed signi cant activation in the bilateral occipital area (cluster-based permutation test, $p < 0.01$; Fig. 3A, left), whereas the Unatt Neutral comparison mainly happened in the postcentral parietal area (cluster-based permutation test, $p < 0.01$; Fig. 3A, right). We further divided targets based on where they were presented (left versus right), and then calculated the corresponding TRF response and the alpha-band power pro les separately. As shown inFig. 3B, the

a relative-time signal instead of absolute-time course (Crosse et al., 2016; Lalor et al., 2006). Therefore, the observed Att-Unatt temporal difference (i.e., 'sequential sampling' or 'sequential processing') in TRF responses primarily supports their temporal latency difference (defined relative to each luminance transient), and does not necessarily indicate an ongoing shifting between the two locations in absolute time within each trial, given the temporal summation of impulse responses across moments. Moreover, the TRF response and classical VEP have been found to be

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J. Neurosci. Methods 164, 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024.

Marshall, T.R., O'Shea, J., Jensen, O., Bergmann, T.O., 2015. Frontal eye fields control attentional modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex. J. Neurosci.: the of cial journal of the Society for Neuroscience 35, 1638–1647. https://doi.org/10.1523/JNEUROSCI.3116-14.2015.

Michalareas, G., Vezoli, J., van Pelt, S., Schoffelen, J.-M., Kennedy, H., Fries, P., 2016. Alpha-beta and gamma rhythms subserve feedback and feedforward in uences among human visual cortical areas. Neuron 89, 384–397. https://doi.org/10.1016/j.neuron.2015.12.018.

Miller, J., Patterson, T., Ulrich, R., 1998. Jackknife-based method for measuring LRP onset latency differences. Psychophysiology 35, 99–115. https://doi.org/10.1111/1469-8986.3510099.

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput. Intell. Neurosci. 1–9, 2011. https://doi.org/10.1155/2011/156869.

Popov, T., Kastner, S., Jensen, O., 2017. FEF-controlled alpha delay activity precedes stimulus-induced gamma-band Activity in visual cortex. J. Neurosci. 37, 4117–4127. https://doi.org/10.1523/JNEUROSCI.3015-16.2017.

Schroeder, C.E., Lakatos, P., 2009. Low-frequency neuronal oscillations as instruments of sensory selection. Trends Neurosci. 32, 9–18. https://doi.org/10.1016/j.tins.2008.09.012.

Sokoliuk, R., Mayhew, S.D., Aquino, K., Wilson, R., Brookes, M.J., Francis, S.T., Hanslmayr, S., Mullinger, K.J., 2018. Two spatially distinct posterior alpha sources ful ll different functional roles in attention. bioRxiv. https://doi.org/10.1101/384065.

Song, K., Meng, M., Chen, L., Zhou, K., Luo, H., 2014. Behavioral oscillations in attention: rhythmic pulses mediated through band. J. Neurosci. 34, 4837–4844. https://doi.org/10.1523/JNEUROSCI.4856-13.2014.

Spaak, E., Bonnefond, M., Maier, A., Leopold, D.A., Jensen, O., 2012. Layer-speci c entrainment of gamma-band neural activity by the alpha rhythm in monkey visual cortex. Curr. Biol. 22, 2313–2318. https://doi.org/10.1016/j.cub.2012.10.020.

Tsotsos, J.K., 1990. Analyzing vision at the complexity level. Behav. Brain Sci. 13, 423–445. https://doi.org/10.1017/S0140525X00079577.

VanRullen, R., Macdonald, J.S.P., 2012. Perceptual echoes at 10 Hz in the human brain. Curr. Biol. 22, 995–999. https://doi.org/10.1016/j.cub.2012.03.050.

Walter, S., Quigley, C., Andersen, S.K., Mueller, M.M., 2012. Effects of overt and covert attention on the steady-state visual evoked potential. Neurosci. Lett. 519, 37–41. https://doi.org/10.1016/j.neulet.2012.05.011.

Wang, Y., Luo, H., 2017. Behavioral oscillation in face priming: prediction about face identity is updated at a theta-band rhythm. Progress in Brain Research. Elsevier, pp. 211–224. https://doi.org/10.1016/bs.pbr.2017.06.008.

Zumer, J.M., Scheeringa, R., Schoffelen, J.-M., Norris, D.G., Jensen, O., 2014. Occipital alpha activity during stimulus processing gates the information ow to object-selective cortex. PLoS Biol. 12, e1001965. https://doi.org/10.1371/journal.pbio.1001965.